arXiv:1208.0618v1 [quant-ph] 2 Aug 2012

Time periodicity and dynamical stability in two-boson systems

Jose Reslen[1]
[1] *Coordinación de Física, Universidad del Atlántico,*
Km. 7 Antigua vía a Puerto Colombia, Barranquilla, Colombia.

(Dated: August 6, 2012)

We calculate the period of recurrence of dynamical systems made of two interacting bosons. A number of theoretical issues related to this problem are discussed, in particular, the conditions for small periodicity. The knowledge gathered in this way is then utilized to propose a notion of dynamical stability based on the stability of the period. Dynamical simulations show good agreement with the proposed scheme.

PACS numbers:

INTRODUCTION

An important result known as the Recurrence Theorem [1, 2] establishes that classical as well as quantum systems of constant energy will come back very close to their initial state at some (maybe long) time during their evolution. A straightforward consequence of the theorem is that these systems posses, at least in a broad sense, an intrinsic or natural frequency given by the inverse of the recurrence's time. Such a frequency characterizes the system's response to external driving. For instance, the amplitude of the response is maximally enhanced when the frequency of the driving matches the natural frequency of the system [3]. This enhancement underlies quite a number of important physical phenomena and its understanding is of fundamental interest. Assuming that the frequency of the driving is constant, one should minimize fluctuations of the natural frequency in order to get an amplified response, especially when the parameters of the system are subject to small perturbations. The stability of quantum dynamics as a result of a small change in the parameters has been discussed in several works, for example in chapter 11 of [4] or [5]. It is seen that in models with classical counterparts, the overlap of the quantum state of two systems with the same initial condition but slightly different parameters remains close to one as long as the initial quantum condition is well localized inside a stable island of the classical map. The opposite takes place when the initial condition is localized in a chaotic region. Here we intend to approach the issue of stability by studying the behavior of periodicity in a model with no direct classical analog. The study contains a moderate analysis of the period, which lays the foundation for the subsequent argument about stability. In absence of a classical analog, we base our approach exclusively on the eigenenergies of the Hamiltonian, making little reference to the state. We discuss a series of issues in connection to the period, particularly, the question of how small the period can be as a function of the parameters. The results obtained here are relevant in the context of quantum computation, where some information protocols or quantum gates are sensitive to perturbations of the parameters and periodic systems can be seen as quantum memories [7].

In the language of the theory of second quantization, the quantum state is written with reference to occupation modes of unperturbed levels. In this context, let us focus on a model featuring two-body interaction such as,

$$\hat{H} = J(\hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1) + \sum_{k=1}^{2} \frac{U_k}{2} \hat{a}_k^\dagger \hat{a}_k (\hat{a}_k^\dagger \hat{a}_k - 1) + \epsilon_k \hat{a}_k^\dagger \hat{a}_k. \quad (1)$$

As usual, the exchange term J and the unperturbed energies ϵ_1 and ϵ_2 underline the single body response while U_1 and U_2 determine the intensity of contact interaction among particles. The mode operators satisfy the usual bosonic relations $[\hat{a}_1, \hat{a}_1^\dagger] = [\hat{a}_2, \hat{a}_2^\dagger] = 1$, etc. The unperturbed system can be probed by looking at the absorption profile of an incident laser of frequency $\nu = |\epsilon_2 - \epsilon_1|/\hbar$. The total number of particles,

$$M = \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_2^\dagger \hat{a}_2, \quad (2)$$

is a preserved quantity. The proposed system reduces to a two-level model when $M = 1$. Hamiltonian (1) can be rearranged as,

$$\hat{H} = \eta \hat{a}_1^\dagger \hat{a}_1 \hat{a}_1^\dagger \hat{a}_1 - \mu \hat{a}_1^\dagger \hat{a}_1 + \hat{a}_1^\dagger \hat{a}_2 + \hat{a}_2^\dagger \hat{a}_1, \quad (3)$$

where, for $M = 2$, the parameters η, μ and Δ turn out to be,

$$\eta = \frac{U_1 + U_2}{2J},\ \mu = \frac{3U_2 + U_1}{2J} + \frac{\Delta}{J},\ \Delta = \epsilon_2 - \epsilon_1. \quad (4)$$

In writing the previous identities we have established J as our energy unit [9]. The state evolution is given by (in what follows, we set $\hbar = 1$),

$$|\psi(t)\rangle = q_1 e^{-iE_1 t}|E_1\rangle + q_2 e^{-iE_2 t}|E_2\rangle + q_3 e^{-iE_3 t}|E_3\rangle. \quad (5)$$

Hence, periodicity arises whenever [8],

$$E_1T = 2\pi n_1,\ E_2T = 2\pi n_2,\ E_3T = 2\pi n_3. \tag{6}$$

Being T the period of the recurrence and n_1, n_2 and n_3 integer numbers. In this case periodicity is absolute as the quantum state recurs identically at regular intervals and the corresponding evolution operator equals the unity operator. Another form of periodicity [6] emerges by considering instances in which the quantum state recurs up to a phase, i.e.,

$$|\psi(t+\tau)\rangle = e^{-i\phi}|\psi(t)\rangle. \tag{7}$$

We refer to this as partial periodicity as the phase factor might generate quantum interference effects. As an example, let us look at Hamiltonian (3) for a single particle,

$$\hat{H} = \begin{pmatrix} -\mu & 1 \\ 1 & 0 \end{pmatrix}. \tag{8}$$

Absolute periodicity occurs whenever the ratio of energies is a fractional number,

$$x = \frac{E_1}{E_2} = \frac{\mu - \sqrt{\mu^2+4}}{\mu + \sqrt{\mu^2+4}} = \frac{n_1}{n_2}. \tag{9}$$

T can be found from Eq. (6),

$$T = 2\pi n_2\sqrt{-x} = 2\pi\sqrt{-n_1n_2}. \tag{10}$$

In Eq. (9) we can define E_1 and E_2 so that $|n_2| \geqslant |n_1|$ and therefore $-1 \leqslant x < 0$. In principle, there is no limit on the maximum value of T. Conversely, the minimum value is $T = 2\pi$ and takes place at $\mu = 0$.

In a similar way, partial periodicity derives from,

$$E_1\tau = \phi,\ E_2\tau = \phi + 2\pi, \tag{11}$$

and therefore,

$$\tau = \frac{2\pi}{|E_2 - E_1|} = \frac{2\pi}{\sqrt{\mu^2+4}} = 2\pi\sqrt{\frac{-x}{(1-x)^2}}. \tag{12}$$

The first equality is consistent with the view that the system's natural frequency is proportional to the difference of its two eigenenergies. Unlike T, τ reaches a maximum of $\tau = \pi$ at $\mu = 0$ and goes down asymptotically to zero as $\mu \to \pm\infty$, where one of the eigenenergies dominates the spectrum and the Hamiltonian is almost singular. This shows that τ is maximum when T is minimum and that τ goes to zero as T goes to infinity. Two level systems always display partial periodicity, but not always absolute periodicity.

TWO PARTICLES

Let us now probe these periodicity concepts in a bigger system. Hamiltonian (3) for $M = 2$ takes the matrix form,

$$\hat{H} = \begin{pmatrix} 4\eta - 2\mu & \sqrt{2} & 0 \\ \sqrt{2} & \eta - \mu & \sqrt{2} \\ 0 & \sqrt{2} & 0 \end{pmatrix}. \tag{13}$$

The energies are the solutions of the characteristic equation,

$$E^3 + \alpha E^2 + \beta E + \gamma = 0, \tag{14}$$

where,

$$\alpha = -5\eta + 3\mu, \tag{15}$$
$$\beta = 2(2\eta - \mu)(\eta - \mu) - 4, \tag{16}$$
$$\gamma = 4(2\eta - \mu). \tag{17}$$

From the previous equalities we can see that if an energy E is a solution of Eq. (14) for a set of parameters $\{\eta, \mu\}$, then $-E$ is a solution for the set $\{-\eta, -\mu\}$. Additionally, given two solutions of Eq. (14), E_3 and E, we can write,

$$E_3^3 - E^3 + \alpha(E_3^2 - E^2) + \beta(E_3 - E) = 0. \tag{18}$$

If $E_3 \neq E$, the polynomial can be simplified as,

$$E^2 + E(E_3 + \alpha) + E_3^2 + \alpha E_3 + \beta = 0, \tag{19}$$

and the respective solutions provide the unaccounted energies,

$$E_1 = -\frac{E_3 + \alpha + \sqrt{(E_3+\alpha)^2 - 4(E_3^2 + \alpha E_3 + \beta)}}{2}, \tag{20}$$

and similarly for E_2. Absolute periodicity results when the energy ratios adhere to,

$$x = \frac{E_1}{E_3} = \frac{n_1}{n_3},\ y = \frac{E_2}{E_3} = \frac{n_2}{n_3}. \tag{21}$$

One may ask whether given a set of parameters η and μ the system would display periodicity. This however might not be a convenient approach, since in any case we can find integers n_1, n_2 and n_3 showing ratios as close to x and y as we want. This is also true for any reasonable quantum closed system. Instead, we propose an approach in which given a pair of ratios we ask if a set of parameters producing these ratios do exist. Following this idea we proceed as follows.

Standard Procedure

Let us then consider x and y as known variables. We divide Eq. (20) by E_3 and get x. y is obtained in a similar way. By means of algebraic operations we can put the unknown variables in terms of α,

$$E_3 = -\alpha/(1+x+y), \tag{22}$$
$$\beta = p\alpha^2, \tag{23}$$
$$\gamma = q\alpha^3. \tag{24}$$

So that p and q are given by,

$$p = (x+y+xy)/(1+x+y)^2, \tag{25}$$
$$q = xy/(1+x+y)^3. \tag{26}$$

From equations (15) and (17) we get,

$$\gamma = \frac{4}{3}(\eta - \alpha). \tag{27}$$

Similarly, using (27) in combination with (24) leads to,

$$\eta = \alpha\left(1 + \frac{3q}{4}\alpha^2\right). \tag{28}$$

Inserting (28) in (15) yields,

$$\mu = 2\alpha\left(1 + \frac{5q}{8}\alpha^2\right). \tag{29}$$

Combining equations (16), (23), (28) and (29) we get the characteristic equation,

$$q^2A^3 + 2qA^2 + 4pA + 16 = 0, \tag{30}$$

where $A = \alpha^2$. In this way, given a pair of values x and y we find p and q from Eqs. (25) and (26) and introduce them in Eq. (30). From the solutions we have α and therefore the η and μ that deliver the energy ratios. Strictly, we have 6 solutions, but they come in pairs giving energies with opposite sign. In order for a solution to be physically acceptable we demand that η and μ should be real. Let us now discuss particular cases where the previous method does not apply.

Particular Cases

Case A. $\alpha = x + y + 1 = 0$.



FIG. 1: Coordinate map indicating the number of set of parameters $\{\eta, \mu\}$ delivering energy ratios $\{x, y\}$. Red (dark gray) region: 0 solutions. Green (light gray) region: 4 solutions. Blue (continuous) lines: 2 solutions. Yellow dots ($\{x = -1, y = 0\}$ and $\{x = 0, y = -1\}$): 1 solution. Black (dashed) lines highlight the pairs for which there are solutions with $\eta = 0$. In these cases, given a couple $\{x, y\}$ there are two solutions with $\eta = 0$ and two solutions with $\eta \neq 0$. The mirror symmetry around $y = -x$ derives from the fact that E_1 and E_2 can be swapped in Eqs. (21). Here we require $|E_1| \leqslant |E_3|$ and $|E_2| \leqslant |E_3|$. Along the edges of the square where x or y is -1, there exists an ambiguity in the choice of E_3 because two eigenenergies display the same absolute value but opposite sign. This implies that the ratio that is not -1 can be positive or negative. As a consequence, the borders remain identified: $\{x = -1, y\} \sim \{x = -1, -y\}$ and $\{y = -1, x\} \sim \{y = -1, -x\}$.

For a pair of values x and y the parameter η is a solution of the polynomial equation (see appendix A),

$$\kappa^2\eta^6 + 27\kappa^2\eta^4 + \left(243\kappa^2 + \frac{81}{4}\right)\eta^2 + 729\kappa^2 = 0, \tag{31}$$

in which we have introduced,

$$\kappa^2 = -\frac{4}{3+(x-y)^2}\left(1 - \frac{4}{3+(x-y)^2}\right). \tag{32}$$

Moreover, μ can be found from Eq. (15).

Case B. $x = 1$ or $y = 1$ (excluding $\{x = 1, y = 0\}$, $\{x = 0, y = 1\}$ and $\{x = y = 1\}$).

In order to avoid division by zero in Eq. (18) we modify the variables in the following fashion,

$$\text{if } x = 1 \Rightarrow x' = y' = \frac{1}{y}, \tag{33}$$

and analogously when $y = 1$. In this way the new variables admit the standard procedure.



FIG. 2: Top: T (arbitrary units) vs. x, keeping y constant. Bottom: T vs. y, keeping x constant. In both cases the grid slice is 1/97. As can be seen, the smallest values of T are found close to the origins.

Case C. $\{x = 0, y = 1\}$ or $\{x = 1, y = 0\}$.

Then $q = \gamma = 0, \beta = -4$ and $\alpha = \eta$. Non-vanishing energy values satisfy $E^2 + \eta E - 4 = 0$ and x (or y) $= 1$ is a solution only if $\eta = \pm 4i$.

Case D. $\{x = y = 1\}$.

The spectrum of the Hamiltonian is three fold degenerate. Hence the eigenvalue equation is of the form $(E_3 - E)^3 = 0$. Comparing with Eq. (14) we infer $-3E = \alpha, 3E^2 = \beta$ and $-E^3 = \gamma$. It then follows $E^6 + 6E^4 + 12E^2 + 16 = 0$, which has no real solutions and therefore no real-valued η and μ produce a fully degenerate spectrum.

Fig. 1 shows a map classifying the coordinate space according to the number of valid solutions encountered for every pair of ratios $\{x, y\}$. The maximum number of solutions is 4, usually coming from 2 real solutions of Eq. (30). Along the negative side of the axes the Hamiltonian becomes reducible with $\mu = 2\eta$, hence only two solutions are possible. The two instances with 1 solution correspond to $\eta = \mu = 0$.

Recurrences

The condition for absolute periodicity is given in Eq. (5) while partial periodicity can be determined through,

$$E_1\tau = \phi, \; E_2\tau = \phi + 2\pi N_2, \; E_3\tau = \phi + 2\pi N_3, \qquad (34)$$

in such a way that N_2 and N_3 are both integers. If $T = N\tau$, being N an integer, we can infer from Eqs. (6) and (34),



FIG. 3: Partial period (arbitrary units) against x for $M = 2$ and $M = 1$ (Eq. (12) - dashed blue line). We have intentionally restrained the vertical axis to $0 < \tau < 10$. Interacting Hamiltonians can display longer τ than non-interacting Hamiltonians. For this particular graph we used grid slices ranging from 1 to 1/500 in order to scan the xy square of Fig 1 (see appendix B). The inclusion of finer grids would produce a denser, but not very different, graph. This is because finer grids are probabilistically related to bigger periods.

$$\phi = \frac{2\pi n_1}{N}, \; N_2 = \frac{n_2 - n_1}{N}, \; N_3 = \frac{n_3 - n_1}{N}. \qquad (35)$$

In order to find T and τ, we first determine n_1, n_2 and n_3 from the rationals x and y in such a way that there is no common divisor greater than 1 among the three generating integers. Simultaneously, the values x and y are used to encounter corresponding eigenenergies following the previously discussed methodology. We are then entitled to employ any of the identities of Eqs. (6) (in particular we employ $E_3 T = 2\pi n_3$) to get T using one of the eigenenergies. Finally, the integer N results as the greatest common divisor of $n_2 - n_1$ and $n_3 - n_1$. As can be seen, large values of the integer n_3 are more likely to produce large T and τ.

Fig. 2 presents sample diagrams of T against x and y. A complete depiction would be quite more intricate. We point out that T decreases as x and y approach zero. This behavior appears to be generic, laying minimum values of T around (or in) the origins. Following T across the line $\{x = -1/n, y = 1/n\}$ for $n = 5, 6, 7$, etc. we find it goes down asymptotically toward $T = \pi\sqrt{2}$ as $n \to \infty$. The same minimum value can be analytically found at $\{x = -1/2, y = 0\}$ and its equivalent. Both instances suggest a relation between minimum T and the Hamiltonian matrix being or becoming singular.

Fig. 3 pictures the intricate relation between τ and x obtained by testing energy ratios over the xy square of Fig. 1 as explained in appendix B. Cooperative systems may display longer partial periods than non-interacting



FIG. 4: Blue lines: set of parameters for which the period is stable under small changes in the direction indicated by the red lines. Inset. τ (arbitrary units) against the smallest of $|x|$ and $|y|$. It follows that as τ goes to zero one of the eigenenergies becomes small in comparison to the others.

systems, therefore periodicity can help identify interacting phases. This behavior is most likely due to the capacity of many-body systems to develop complex dynamics on account of the increased number of effective states. Also of interest is the chaotic aspect of Fig. 3, meaning that small intervals in xy space do not necessarily map onto small intervals in τ space. This lack of continuity, which also characterizes T, takes place because neighboring values of x and y do not always correspond to neighboring values of n_1, n_2 and n_3. For instance, $\{x = 1/3, y = 2/3\}$ is not far from $\{x = 97/300, y = 201/300\}$, but the sets of integers that generate each pair are widely different. Recall that the period depends directly on the integers. In addition, there can be common divisors between $n_2 - n_1$ and $n_3 - n_1$ and this might further affect the continuity of τ. Finally, it can be seen from the inset of Fig. 4 that when τ becomes small at least one of the ratios x or y approaches zero, once again suggesting that in a Hamiltonian displaying small periodicity one of the eigenenergies is much smaller than the others.

STABILITY

As introduced here, periodicity, either total or partial, is a characteristic of the Hamiltonian. It does not depend (except when the initial state is an eigenstate of the Hamiltonian) on the state. This encourage us to ask whether one can change the Hamiltonian parameters without affecting the period. This is what happens for one particle when $\mu = 0$, as can be seen from Eq. (12):

$$\left(\frac{d\tau}{d\mu}\right)_{\mu=0} = 0. \tag{36}$$

For two particles, we can start by arguing that the period is stable whenever both x and y are stable under changes of the parameters. Moreover, we can consider the energy ratios as functions of the parameters, $x(\eta, \mu)$ and $y(\eta, \mu)$, in such a way that maximum variations occur in the direction of the function gradients and zero variations take place in the directions perpendicular to the gradients. In this way, for any set of parameters η and μ one can always identify a direction of change of the parameters along which *one* of the ratios is stable. It then follows that in order for T to be stable the directions of zero change of both ratios must coincide, i.e., the gradients must be parallel:

$$\nabla x + \lambda \nabla y = 0, \tag{37}$$

where $\nabla x = \frac{\partial x}{\partial \eta}\vec{\eta} + \frac{\partial x}{\partial \mu}\vec{\mu}$ and similarly for ∇y. $\vec{\eta}$ and $\vec{\mu}$ are unitary vectors in the parameter space and λ is a real number. This problem is equivalent to finding the extreme values of $x(\eta, \mu)$ subject to the condition $y(\eta, \mu)$=constant, or the other way around. In this context λ takes the roll of a Lagrange multiplier and the analogy applies as long as the involved functions are smooth. The extremes of x can be worked out from Fig. 1. It is found, however, that not all extreme values imply parallel gradients. For instance, along the edges, where x or y is -1, the identification of borders on each side of the axes generates a discontinuity in the first derivative of the ratios. In addition, for the extremes located around the neighborhood of the axes at least one of the parameters diverge toward infinity. Let us remark that there is no solution of Eq. (30) along the positive side of the axes. In general, we found the gradients are parallel only along the blue lines of Fig. 1 located in between green and red regions.

Fig. 4 shows curves indicating the parameters as well as the directions of change corresponding to zero variation of the period. Far from the origin the curves approach (but do not seem to touch) straight lines given by simple expressions. In one case the direction of zero change aligns with the direction of the curve as we get away from the origin. In the other case the direction of zero change becomes constant with an angle of inclination θ satisfying $\tan\theta \approx 1/\sqrt{2}$.

As can be seen from Fig. 5, when the parameters are shifted in the direction of zero change the dynamics of corresponding systems look similar. Conversely, when the change of parameters occurs in the direction of the gradient, the dynamics soon diverge. This behavior is consistent with our study and shows that in some cases the period alone characterizes the global profile of the system evolution.

We have studied time periodicity as well stability properties of interacting bosonic systems. We stressed the difference between the usual notion of periodicity, where



FIG. 5: Mean number of particles $\langle \hat{a}_1^\dagger \hat{a}_1 \rangle$ against time (arbitrary units) for three slightly different cases. Top. $\eta = 1.878615$ and $\mu = 2.939702$. Middle. $\eta = 1.778615$, $\mu = 2.775950$. Bottom. $\eta = 1.878615$ and $\mu = 2.714882$. The middle-graph corresponds to a set of parameters in the curve of stability of Fig. 4. Top- and bottom-graphs show the evolution of the system when such parameters are shifted in the direction of zero change and the direction of the gradient, respectively.

the ratios of the Hamiltonian energies become commensurate, and partial periodicity, where the state recurs up to a phase factor. Both forms of periodicity were explicitly established for one and two particles in a model described by two bosonic modes. The results suggest a connection between minimum periodicity and the fact that one of the eigenvalues becomes small against the other eigenvalues, especially for the two-particle case [10]. Likewise, we observed that the stability of the period does not only depend on the parameters but also on the direction of change of such parameters. Hence, we presented a diagram showing the set of parameters as well as the directions of change for which the period stays constant. These results were consistent with simulations of the dynamics.

An issue of the method previously applied to find τ is the assumption $T = N\tau$ for the two-particle case. It may be possible to relax this assumption and derive τ using only Eq. (34). This would in principle lead to a richer stability diagram including parameters for which τ is stable but T is not, e.g., $\eta = \mu = 0$. Similarly, we feel that the method presented here is robust, although slightly dependent on the dimension of the Hamiltonian.

Appendix A: Derivation of Eq. (31).

When $\alpha = 0$ the solutions of Eq. (19) can be written as,

$$E_1 = \frac{-E_3 - \sqrt{-3E_3^2 - 4\beta}}{2}, \quad E_2 = \frac{-E_3 + \sqrt{-3E_3^2 - 4\beta}}{2}. \tag{38}$$

Then it follows,

$$E_3 = \sqrt{\frac{-4\beta}{3 + (x-y)^2}}. \tag{39}$$

Replacing (39) in (14) with $\alpha = 0$ gives,

$$\kappa \beta^{3/2} = \gamma, \tag{40}$$

where κ is given by Eq. (32). Likewise, $\mu = \frac{5}{3}\eta$. By replacing such a value of μ in Eqs. (16) and (17) we find,

$$\beta = -4\left(\frac{\eta^2}{9} + 1\right), \quad \gamma = \frac{4}{3}\eta. \tag{41}$$

Substituting these values in Eq. (40) and after some algebraic steps we arrive at Eq. (31).

Appendix B: Generation of commensurate ratios.

In order to generate pairs of ratios compatible with the coordinate square of Fig. 1, we first establish an integer $n_3 = 1, 2, 3, \ldots$, which is also the inverse of the grid slice. For a given n_3, we check $-n_3 < n_1 < -1$ and $n_1 < n_2 < n_3 - 1$. The values of n_3 are introduced into a computer routine in increasing order starting from $n_3 = 1$. Subsequently, n_1 and n_2 are introduced. The integers are used to get x and y which are in turn used to check for Hamiltonians with corresponding eigenenergies. As the coordinate square is symmetric, we scan only the section $x < 0$. If for a given triplet $\{n_1, n_2, n_3\}$ there exists a maximum common divisor greater than 1, the triplet is discarded. Notice that in this way the scan of the coordinate square is performed using several superimposed grids, avoiding repetition of pairs $\{x, y\}$.

[1] L. Barreira, *XIVth International Congress on Mathematical Physics*, 415-422 (2006).
[2] D.L. Shepelyansky, *Phys. Rev. E* **82**, 055202(R) (2010).
[3] J. Reslen, C.E. Creffield and T.S. Monteiro, *Phys. Rev. A* **77**, 043621 (2008).
[4] A. Peres, *Quantum Theory: Concepts and Methods*, ed. by A.V.D Merwe, Kluwer Academic Publishers, (2002).
[5] G. Casati and T. Prosen, *Braz. J. Phys.* **35**, 233 (2005).
[6] P.P. Rohde, A. Fedrizzi, T.C. Ralph, *J. Mod. Opt.* **59**, 710 (2012).
[7] S.M. Giampaolo, F. Illuminati, A. Di Lisi, G. Mazzarella, *Int. J. Quant. Inf.* **4**, 507 (2006).

[8] C. Godsil, *Elec. J. Comb.*, **18** 23 (2011).
[9] As J usually depends on the details of the system, we avoid making reference to it and use arbitrary units for our time scales.
[10] The only exception takes place for one particle when $\mu \to \infty$ since according to Eqs. (9) and (10) $T \to \infty$.